

CASE REVIEW

Infrapopliteal Thromboembolic Arterial Occlusion

Case Demographics

Age (years): 75
Gender: Male

Clot Removal

Pounce™ Thrombectomy System
(no thrombolysis or aspiration)

Clinical Presentation


- Pain and paresthesia of right foot
- Symptom duration 1 day
- Dx of new-onset atrial fibrillation
- DUS evidence of thromboembolism in right lower extremity
- Angiography revealed 100% thrombotic occlusion of multiple below-the-knee vessels

Procedural Information


The patient received three passes of the Pounce™ Thrombectomy System to remove thromboembolic clot from the right anterior tibial, posterior tibial, and peroneal arteries, with balloon angioplasty in the posterior tibial. Post-procedure ABI was 1.1 with normal Doppler velocities. No thrombolysis used.

Anatomy

Pre



Post



INTENDED USE: The Pounce™ Thrombectomy System is intended for the non-surgical removal of thrombi and emboli from the peripheral arterial vasculature.

CONTRAINDICATIONS:

- The device is contraindicated for use in patients who cannot receive recommended intravenous anticoagulant therapy.
- The device is not for coronary or neurovascular use.
- The device is not intended to be deployed in vessels with previously implanted devices.

Refer to Instructions for Use for more information concerning Indications, Contraindications, Specific Anatomic Considerations, Warnings, Precautions, and Adverse Events.

CAUTION: Federal (USA) law restricts this device to sale by or on the order of a physician.

NOTE: Not all product components are available in every country.

Please consult with your Endologix representative to confirm product availability.

Endologix, Pounce and the Endologix and Pounce logos are trademarks of Endologix LLC and/or its affiliates. All other trademarks are the respective property of their owners.

© 2026 Endologix LLC. All rights reserved. MM3063-US Rev 01